

TAXON TABLE FOR SANDERSON AND DOYLE, 2001

18S Species (1)	Source (2)	rbcL Species	Source
<i>Chara vulgaris</i>	U81271	<i>C. connivens</i>	L13476
<i>Marchantia polymorpha</i>	X75521	...	X04465
<i>Lycopodium phlegmarium</i> = <i>Huperzia phlegmaria</i>	X81964	<i>L. digitatum</i>	L11055
<i>Equisetum robustum</i>	X78890	<i>E. arvense</i>	L11053
<i>Osmunda cinnamomea</i>	U18516	...	D14882
<i>Asplenium nidus</i> = <i>A.</i> <i>australasicum</i>	D85303	...	U05907
<i>Cycas taitungensis</i>	D85297	<i>C. circinalis</i>	MC
<i>Zamia pumila</i>	M20017	<i>Z. inermis</i>	MC
<i>Ginkgo biloba</i>	D16448	...	MC
<i>Pinus elliotii</i>	D38245	<i>P. radiata</i>	MC
<i>Nageia nagi</i>	D16447	<i>Podocarpus gracilior</i>	MC
<i>Taxus mairei</i>	D16445	<i>T. brevifolia</i>	MC
<i>Ephedra torreyana</i>		<i>E. tweediana</i>	MC
<i>Welwitschia mirabilis</i>	D85299	...	MC
<i>Gnetum gnemon</i>	DS	...	MC
<i>Nymphaea tuberosa</i>	DS	<i>N. odorata</i>	MC
<i>Saururus cernuus</i>	DS	...	MC
<i>Austrobaileya scandens</i>	DS	...	MC
<i>Chloranthus spicatus</i>	DS	<i>C. japonicus</i>	MC
<i>Magnolia acuminata</i>	DS	<i>M. sp.</i>	MC
<i>Eupomatia sp.</i>	DS pers. comm.	<i>E. bennettii</i>	MC
<i>Calycanthus floridus</i>	DS	...	L14291
<i>Sassafras albidum</i>	DS	<i>Persea americana</i>	MC
<i>Drimys aromatica</i>	DS	<i>D. winteri</i>	MC
<i>Calla palustris</i>	DS	<i>Spathiphyllum wallisii</i>	MC
<i>Tacca sp.</i>	DS	<i>T. chantrieri</i>	MC
<i>Veitchia sessilifolia</i>	DS	<i>Drymophloeus subdistichus</i>	MC
<i>Oryza sativa</i>	DS	...	MC
<i>Ranunculus sardous</i>	DS	<i>T. trichophyllus</i>	MC
<i>Nelumbo lutea</i>	DS	...	MC
<i>Platanus occidentalis</i>	DS	...	MC
<i>Buxus sempervirens</i>	DS	<i>Pachysandra procumbens</i>	MC
<i>Pisum sativum</i>	DS	...	MC
<i>Fagus sp.</i>	DS. pers. comm.	<i>F. americanus</i>	MC
<i>Carya sp.</i>	DS pers. comm.	<i>C. glabra</i>	MC

<i>Arctostaphylos uva-ursi</i>	DS	<i>Enkianthus campanulatus</i>	MC
<i>Brunfelsia pauciflora</i>	DS	<i>Nicotiana tabacum</i>	MC

Notes

- (1) Species used to represent the same terminal taxon are found in the same row. If no entry is present under rbcL, then the same species was used for both genes.
- (2) Key to sources: DS = Soltis et al., 1997 and Chaw et al. 1997, as supplied by Doug Soltis, pers. comm.; MC = Chase et al. 1993, as supplied by Michael J. Donoghue, pers. comm.; numbers are GenBank accession numbers for other taxa not contained in those analyses; three sequences were supplied by Doug Soltis, as pers. comm. but not contained in published studies